

ORIGINAL RESEARCH



Using leaf economic spectrum and photosynthetic acclimation to evaluate the potential performance of wintersweet under future climate conditions

Lei Wang^{1,2} | Qing-Lai Dang²

¹Department of Landscape Architecture, Jiyang College, Zhejiang A&F University, Zhejiang, China

²Faculty of Natural Resources Management, Lakehead University, Thunder Bay, Ontario, Canada

Correspondence

Lei Wang,
Email: rockyking@163.com

Qing-Lai Dang,
Email: qdang@lakeheadu.ca

Funding information

Natural Sciences and Engineering Research Council of Canada, Grant/Award Number: DDG-2020-00008; Zhejiang A and F University, Grant/Award Number: RC2023B06

Edited by C.H. Foyer

Abstract

The function of landscape plants on the ecosystem can alleviate environmental issues of urbanization and global change. Global changes due to elevated CO₂ affect plant growth and survival, but there is a lack of quantitative methods to evaluate the adaptability of landscape plants to future climate conditions. Leaf traits characterized by leaf economic spectrum (LES) are the universal currency for predicting the impact on plant ecosystem functions. Elevated CO₂ usually leads to photosynthetic acclimation (PC), characterised by decreased photosynthetic capacity. Here, we proposed a theoretical and practical framework for the use of LES and PC to project the potential performance of landscape plants under future climatic conditions through principal component analysis, structural equation modelling, photosynthetic restriction analysis and nitrogen allocation analysis. We used wintersweet (an important landscaping species) to test the feasibility of this framework under elevated CO₂ and different nitrogen (N) supplies. We found that elevated CO₂ decreased the specific leaf area but increased leaf N concentration. The results suggest wintersweet may be characterized by an LES with high leaf construction costs, low photosynthetic return, and robust stress resistance. Elevated CO₂ reduced photosynthetic capacity and stomatal conductance but increased photosynthetic rate and leaf area. These positive physiological traits, e.g., larger leaf area (canopy), higher water use efficiency and stress resistance, may lead to improved performance of wintersweet under the predicted future climatic conditions. The results suggest planting more wintersweet in urban landscaping may be an effective adaptive strategy to climate change.

1 | INTRODUCTION

Continued urbanization and climate change are major challenges humans face in this century (IPCC, 2022; Dervishi et al., 2023). Urban vegetation is an important green infrastructure and is considered key to mitigating these adverse challenges (Velasco et al., 2016; Sousa-Silva et al., 2023). This is based on the biological traits of landscape plants, especially woody plants (Gratani and Varone, 2007a; Liu and Slik, 2022). Canopy shading and transpiration can alleviate the heat island effect caused by urbanization and improve people's thermal comfort (Dujardin

et al., 2022). Plants can reduce the magnitude of climate change by fixing atmospheric CO₂ in photosynthesis and improving air quality (Kim et al., 2020). The seasonal changes in leaf colour, flowers and fruits of plants enhance the natural experience and promote the aesthetic, as well as improve the mental health and increase the age of the residents (Stas et al., 2020; Dujardin et al., 2022; Tang et al., 2023). Landscape vegetation can also reduce the surface runoff of stormwater and noise pollution as well as improve microclimate conditions (Bae and Ryu, 2015; Stas et al., 2020; Downtin et al., 2023). and improve people's health and well-being (Dujardin et al., 2022; Tang et al., 2023).

Due to the unique and irreplaceable ecosystem service and functions, increasing the planting of landscape plants is encouraged in urban planning around the world (Nyelele and Kroll, 2021; Dervishi et al., 2023). It should be noted that poor health conditions of landscape plants can substantially reduce the magnitude of their services and functions (Simon et al., 2018; Esperon-Rodriguez et al., 2023). However, the options of plant species for landscaping are limited, and some species require higher maintenance and management costs than others (Lo Piccolo et al., 2023). Therefore, urban landscape planning requires a wealth of expertise (Ordóñez Barona et al., 2022). Only with a deep understanding of the environmental conditions of the planting site, the ecological requirements and physiological traits of different landscape tree species, and the needs of the community can the scientific species composition and the sustainability of green landscaping be achieved (Alizadeh and Hitchmough, 2020; Nyelele and Kroll, 2021). However, the eco-physiological performance of landscaping plant species in the future under the predicted new environmental conditions is generally poorly understood (Nitschke et al., 2017; Killi et al., 2018). Global change can affect the physiology, growth and survival of plants, but relevant research generally focuses on forest tree species and crops (Nitschke et al., 2017; Kim et al., 2020). The impact of climate change on the functions of urban landscape plants is rarely assessed (Falster et al., 2018; Esperon-Rodriguez et al., 2023). This lack of knowledge seriously restricts the development of proper adaptive strategies in urban landscape planning and predicting landscape plant performance under future climate conditions (Killi et al., 2018).

There is a coordinated relationship between leaf traits and plant functions, which can be used to predict plant responses to global changes (Cui et al., 2020). This is because leaf traits reflect the optimal phenotype the growing environment selects for the given range of resources and environmental conditions (Midolo et al., 2019). The leaf traits, including specific leaf area (SLA, ratio of unit leaf area to mass), leaf nitrogen concentration (N_{area}), and net photosynthetic rate (A_n) are known as leaf economic spectra (LES), which represent a broad trade-off between leaf structure and physiological and ecological traits (Onoda et al., 2017; Yang et al., 2019). SLA relates to leaf thickness and tissue density and affects the photosynthetic return on resource investment for leaf construction cost (Xiong and Flexas, 2018; Maenpuen et al., 2022). SLA is an important functional trait in LES and for plant growth analysis. Leaf traits are frequently measured in ecology to provide a ‘common currency’ for predicting how global change impacts plant ecosystem function and investment strategies for growth and survival (Firn et al., 2019; Lu et al., 2020).

Elevated atmospheric CO₂ is the foundation and driving force of the escalating global change (Wang et al., 2021). Due to intensive human activities and carbon emissions from the combustion of fossil fuels, CO₂ concentrations in urban areas are significantly higher than in natural environments (Gratani and Varone, 2007b). Recent studies have found that while all landscape plant species fix CO₂ via photosynthesis, the photosynthetic capacity (PC: maximum carboxylation rate (V_{cmax}) and maximum electron transfer rate (J_{max})) varies as much as 2–8 times between different tree species used in landscaping

(Kimm and Ryu, 2015; Lo Piccolo et al., 2023). However, the effects of elevated CO₂ on the physiology and growth of landscape plants are rarely reported. The limited literature suggests that elevated CO₂ can change the growth rate of ornamental plants (Kim et al., 2020). The plants growing under emulated future CO₂ usually exhibit typical photosynthetic acclimation, such as decreased PC and stomatal conductance (g_s) (Knauer et al., 2019; Pao et al., 2019; Smith and Keenan, 2020). As a substrate of photosynthesis, elevated CO₂ generally increases carbohydrate and biomass production. Without a corresponding increase in nitrogen supply and uptake, such an increase will lead to a decrease in nitrogen concentration in the plant, particularly in the leaf, triggering a decline in photosynthetic capacity, i.e., photosynthetic acclimation (Silva-Pérez et al., 2020; Ainsworth and Long, 2021; Krämer et al., 2022). Thus, proper assessment of photosynthetic acclimation to elevated CO₂ may be key for predicting plant response to global change.

Wintersweet (*Chimonanthus praecox* L.) is widely used in landscaping in temperate regions (Shang et al., 2020). It blooms in winter with a sweet fragrance and has high ornamental value. Wintersweet has a unique and irreplaceable position in urban parks due to its unique timing of flowering and fresh, sweet smell (Tian et al., 2019a; Song and Wu, 2022). It is often used by landscape architects as an ornamental highlight for landscape routes and an esthetic attraction to visitors in the winter (Shen et al., 2021). Past studies of wintersweet mainly focus on its floral fragrance and drought resistance (Anirban, 2019; Tian et al., 2019b), and its acclimation to future climate conditions, such as elevated CO₂, has not been reported. Enrichment of CO₂ typically promotes plant biomass, which dilutes the leaf nitrogen concentration in leaves and, in turn, affects photosynthetic capacity and nitrogen supply, influencing the leaf traits in the LES (Ainsworth and Long, 2021). This study investigated the responses of the LES and photosynthetic traits to elevated CO₂ and variations in nitrogen supply under controlled environmental conditions. The results of this study provide a theoretical and practical framework for quantifying and predicting the physiological and growth performance of landscape plants under the predicted future climate conditions and for landscape planning. Our methodologies should also be useful for evaluating the potential performance of other landscape woody species under the predicted future climate conditions.

2 | MATERIALS AND METHODS

2.1 | Plant materials and treatments

The seeds of wintersweet were collected from the landscape arboretum at the Zhejiang A&F University campus (30.2374 N, 119.7284E). The seeds were stratified in wet sand at 4°C for about 60 days before being sown in germination trays. Seedlings were transplanted into 4 L plastic pots filled with vermiculite and peat moss (1:1, v:v) (Sun Gro®).

The experiment was conducted in four phytotrons on the Thunder Bay campus of Lakehead University. The experiment followed a split-plot design with two [CO₂] levels (ambient 400 μmol mol⁻¹

(aCO₂) and elevated 800 μmol mol⁻¹ (eCO₂) as the main plot, and three nitrogen levels as the split-plot (containing 3 mM (N3), 6 mM (N6), and 9 mM (N9) nitrogen). Each CO₂ level had two replicates (two separate phytotrons). Each replicate of each treatment combination had 8 seedlings (a total of 96 seedlings: 2 CO₂ * 2 replicates * 3 N forms * 8 seedlings).

The nitrogen treatments used ammonium nitrate (BioBasic Inc.). All N treatments received identical concentrations of other nutrient elements: 5 mM Potassium phosphate and 4 g L⁻¹ Micromax Micronutrients Granular (calcium 6%, magnesium 3%, sulphur 12%, boron 0.1%, copper 1%, iron 17%, manganese 2.5%, molybdenum 0.05%, zinc 1%) (Everris NA Inc.). The seedlings were fertilized twice weekly and irrigated with 500 mL water every other day. The positions of the seedlings were randomly changed within the same phytotron and between the replicate phytotrons of each CO₂ treatment every two weeks.

The CO₂ concentration in each phytotron was maintained using a CO₂ generator (GEN-2E, Custom Automated Products Inc.). Other environmental conditions in all the phytotrons were 25/16°C (day/night) temperature, 50% RH and 16-h photoperiod. High-pressure sodium lamps (P.L. Systems) were used to supplement the natural light when the ambient light intensity was below 500 μmol m⁻² s⁻¹ or the natural day length was shorter than 16-h. All the environmental variables in each phytotron were monitored and controlled by an Argus Titan Environment system (Argus Control Systems Ltd.).

2.2 | Gas exchange and pigment measurement

Three seedlings were randomly selected from each treatment combination, and foliar gas exchange was measured between 9:00 and 16:00 on an unshaded mature leaf near the top of the canopy using a LI-6800 Portable Photosynthesis System (LI-COR Biosciences). The gas exchange under treatment CO₂ was measured at the corresponding treatment [CO₂] (aCO₂ at 400 μmol mol⁻¹, eCO₂ at 800 μmol mol⁻¹), 25°C temperature, 1.1–1.3 kPa water vapour pressure deficit, 1000 μmol m⁻² s⁻¹ photosynthetically active radiation flux density. At least 30 min was allowed before a steady state reading was taken. The net photosynthetic rate (A_{n-g}), stomatal conductance (g_s), intercellular [CO₂] to ambient [CO₂] ratio (C_i/C_a), and photosynthetic nitrogen uses efficiency ($PNUE = A_{n-g}/N_{area}$) were subjected to statistical analyses. Instantaneous photosynthetic water use efficiency ($iWUE = A_{n-g}/E$, where E is the corresponding transpiration rate).

Photosynthetic response to intercellular [CO₂] (A/C_i) curves were measured at 400, 300, 200, 100, 50, 400, 500, 600, 800, 1000, 1200, 1500 μmol mol⁻¹ [CO₂] at the same temperature, VPD, and PAR as given previously. The maximum rate of Rubisco carboxylation V_{max} and maximum rate of photosynthetic electron transport J_{max} were estimated using the fitaci function, and the transition point (C_{i-t} , A_{n-t}) of biochemical limitation from Rubisco to RuBP regeneration was estimated using findCiTranstion function in the Plantecophys R package from A/C_i data (Duursma, 2015). The initial slope of the A/C_i curve was used as the apparent carboxylation efficiency (ACE).

The initial portion of three A/C_i curves was measured under 200, 150, 100, and 50 μmol mol⁻¹ of [CO₂], and 300, 150, and 75 μmol m⁻² s⁻¹ PAR, and used to estimate the daytime respiratory rate (R_d) and intercellular CO₂ compensation point (C_i^*) using the Walker's slope intercept method (Walker et al., 2016). The variable J method was employed to calculate mesophyll conductance (g_m) (Harley et al., 1992), using the equation $\Gamma^* = C_i^* + R_d / g_m$ (Walker et al., 2016), where Γ^* is the CO₂ compensation point in the absence of day respiration.

After the gas exchange measurement, the sample leaves were immediately removed and stored in a refrigerator (-35°C). Leaf pigment was extracted using 80% acetone, and the 645 and 663 nm absorbance was measured. The leaf chlorophyll concentration (Chl) was calculated with $Chl = 20.2 * A_{645} + 8.02 * A_{663}$, where A_{645} and A_{663} are the absorbance at 645 nm and 663 nm, respectively (Wellburn, 1994).

2.3 | Relative photosynthetic limitation and nitrogen partitioning

The relative limitations of photosynthesis by biochemical (l_b), mesophyll conductance (l_m) and stomatal conductance (l_s) limitations were calculated according to Elferjani (Elferjani et al., 2021): $l_b = g_t / (g_t + \partial A_n / \partial C_c)$; $l_s = (g_t / g_s * \partial A_n / \partial C_c) / (g_t + \partial A_n / \partial C_c)$; $l_m = (g_t / g_m * \partial A_n / \partial C_c) / (g_t + \partial A_n / \partial C_c)$, where g_t is the total conductivity to CO₂ diffusion ($g_t = 1 / (1 / g_m + 1 / g_s)$), $\partial A_n / \partial C_c$ is the initial slope of the A_n to C_c response curve and was calculated using the equation $\partial A_n / \partial C_c = V_{cmax} / (\Gamma^* + K_m)$ (Farquhar et al., 1980), where Γ^* was as noted previously and K_m was calculated by $K_m = K_c (1 + O / K_o)$ (Bernacchi et al., 2001). Stomatal limitation (l_s) was calculated using $(C_a - C_i)/C_a$, C_a and C_i are CO₂ concentrations around the leaf and intercellular space, respectively. The partitioning of leaf nitrogen (N) into carboxylation (N_c), electron transfer (N_e) and light capture (N_v) was calculated using the Trouwborst method (Trouwborst et al., 2011).

2.4 | Biomass and nitrogen measurement

After all the above measurements, the height of all the seedlings was measured. Leaf area was measured using a Regent WinFolia system (Regent Instruments Inc.). The seedlings were then harvested and separated into roots, stems, and leaves, and oven-dried at 75°C for 48 h to constant weight to determine the biomass of different organs and SLA. Area-based leaf N concentration (N_{area}) and carbon concentration (C) were measured using the dry combustion method on a CNS-2000 (LECO Corp.) at the Lakehead University Centre for Analytical Services.

2.5 | Statistical analysis

The effects of CO₂ and N levels on photosynthetic and growth traits of individual seedlings were investigated using a two-way analysis of

variance using the agricolae package in R 4.2.4. A three-way ANOVA was performed on A_{n-g} and $PNUE$. Logarithmic or power transformation was used to transform variables that did not meet the normality and homogeneity assumptions before the ANOVA. When an interaction or N effect was statistically significant, LSD post hoc pairwise comparisons were conducted for the means. The PCA function of the FactoMineR package was used for principal component analysis (PCA) to investigate the relationship and clustering of physiological and growth-related parameters. The Bowen method (Bowen et al., 2017) was used to investigate the direct/indirect effects of CO_2 and N levels as categorical variables on biomass and A_{n-g} using the structural equation model (SEM) and the psem function in the piecewiseSEM package (Lefcheck, 2016). Based on the LES and a prior structural model (Onoda et al., 2017; Liu et al., 2022), biomass was explained by

A_{n-g} , N_{area} and SLA (Figure 8a), and A_{n-g} was explained by photosynthetic capacity, Chl and g_t (Figure 8b). The photosynthetic capacity was obtained by the regression equations from V_{cmax} and J_{max} .

3 | RESULTS

3.1 | Biomass and leaf area increased under elevated CO_2 and N enrichment

Elevated CO_2 and increased N supply promoted biomass and total leaf area production in wintersweet, and the values in the eN9 treatment combination were substantially higher than in any other treatments (Figure 1, Table 1). In contrast, e CO_2 significantly reduced SLA ; the




FIGURE 1 Effects of CO_2 and N levels on biomass (a), root shoot ratio (b), photosynthetic rate at growth CO_2 (A_{n-g} , c) and photosynthetic nitrogen use efficiency ($PNUE$, d) in wintersweet. “a-” means ambient CO_2 ($400 \mu mol mol^{-1}$) and “e-” means elevated CO_2 ($800 \mu mol mol^{-1}$) treatments. N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. Means (\pm SE, $n = 6$) with different letters indicated significant differences between treatments (Tukey host hoc test, $P < 0.05$).

TABLE 1 Leaf morphological and photosynthetic traits, and nitrogen concentration of wintersweet grown at ambient and elevated CO_2 response different N levels.

CO_2	N	Leaf area (cm^2)	SLA ($cm^2 g^{-1}$)	N_{area} ($g m^{-2}$)	C/N	L_s	$iWUE$
a CO_2	N3	483 \pm 31 c	143.6 \pm 7.6 a	2.1 \pm 0.1 c	14.1 \pm 0.6 b	0.46 \pm 0.02 bc	7.4 \pm 0.3 ab
	N6	570 \pm 26 c	143.3 \pm 3.9 a	2.3 \pm 0.1 bc	13.4 \pm 0.3 bc	0.58 \pm 0.04 a	7.4 \pm 0.4 ab
	N9	861 \pm 37 b	135.5 \pm 5.2 a	2.6 \pm 0.1 b	12.3 \pm 0.2 c	0.53 \pm 0.04 ab	7.9 \pm 0.6 a
e CO_2	N3	463 \pm 31 c	90.7 \pm 4.4 c	2.3 \pm 0.1 bc	20.1 \pm 0.6 a	0.43 \pm 0.04 cd	6.7 \pm 0.4 ab
	N6	1024 \pm 54 b	120.1 \pm 3.8 b	2.7 \pm 0.1 b	13.9 \pm 0.4 b	0.35 \pm 0.04 d	6.5 \pm 0.4 b
	N9	1325 \pm 91 a	105.8 \pm 4.1 b	3.2 \pm 0.1 a	13 \pm 0.1 bc	0.41 \pm 0.03 cd	6.7 \pm 0.6 ab
P-value	CO_2	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	0.021
	N	< 0.001	0.018	< 0.001	< 0.001	0.7	0.73
	CO_2 -N	< 0.001	0.015	0.24	< 0.001	0.03	0.83

Note: each value represents mean \pm SE ($n = 6$). Two-way ANOVA was performed on the effects of CO_2 , nitrogen (N), and their interaction. Significant effects ($P \leq 0.05$) are shown in bold. Different letters within the same column indicated statistically significant differences between treatments (Tukey post hoc test, $P < 0.05$). a CO_2 : ambient CO_2 ($400 \mu mol mol^{-1}$); e CO_2 : elevated CO_2 ($800 \mu mol mol^{-1}$); N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. SLA : specific leaf area; N_{area} : leaf N content based on leaf-area; C/N: leaf carbon-nitrogen ratio; L_s : stomatal limitation; $iWUE$: intrinsic water use efficiency.

effect was the largest at N3 (Table 1). Furthermore, wintersweet tended to allocate more biomass to the root system under N3 than the other two N treatments (Figure 1b).

3.2 | Leaf N centration, N use efficiency and N distribution

N_{area} increased with the increase in nitrogen supply and CO_2 elevation (Table 1). Elevated CO_2 generally increased, but increasing N supply decreased, the leaf C/N ratio, although not all the effects were statistically significant (Table 1).

Increases in N supply significantly decreased PNUE under a CO_2 but had no significant effect under e CO_2 (Figure 1d). The e CO_2 reduced N allocation to the photochemical apparatus of photosynthesis (Figure 2). The effect of decreasing N supply on N allocation to the biochemical machinery of photosynthesis, however, varied with CO_2 treatment, ranging from having no effects under a CO_2 and decreasing it under e CO_2 (Figure 2).

3.3 | Photosynthetic traits and limitations to photosynthesis

e CO_2 significantly increased A_{n-g} in all the N treatments, but the highest N treatment (N9) significantly increased A_{n-g} only under e CO_2 (Figure 1b). The photosynthesis of wintersweet was limited the most by stomatal resistance (more than 50%) and least by mesophyll resistance (Figure 3). Treatment effects on the traits of A/Ci curves were demonstrated in Figure 4.

Neither CO_2 nor N supply significantly affected leaf chlorophyll concentration (Table 2). The e CO_2 treatment generally reduced g_m




FIGURE 2 Leaf N partitioning into carboxylation (N_c), electron transfer (N_j), and light capture systems (N_v) in response to CO_2 and different N levels in wintersweet. “a-” means ambient CO_2 ($400 \mu\text{mol mol}^{-1}$) and “e-” means elevated CO_2 ($800 \mu\text{mol mol}^{-1}$) treatments. N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively.

and g_s , but the effects were not statistically significant in all the N treatments (Table 2). e CO_2 , however, significantly increased C_i/C_a in all the N treatments (Table 2).

3.4 | Acclimation to CO_2 and N supply

PCA results showed that seedlings grown under a CO_2 converged on the ellipse in the directions of L_s , SLA , $iWUE$ and $PNUE$ (on the left of Figure 5a), while seedlings grown under e CO_2 converged on the A_{n-t} , N_{area} , biomass, A_{n-g} , C/N and C_{i-t} ellipse (on the right of Figure 5a). L_s and SLA on the left were inversely proportional to C_{i-t} , C_i/C_a , and C_{i-g} on the right (Figure 5). With increases in N supply, corresponding ellipses rotated counterclockwise upward (Figure 5b). The seedlings grown under low N supply tended to maintain C/N, while those in high nitrogen treatments were characterized by high values in J_{max} , A_{max} , A_{n-t} , N_{area} , biomass, and A_{n-g} (Figure 5b).

4 | DISCUSSION

4.1 | Leaf economic spectrum and leaf traits

The LES mainly involves SLA , N_{area} , PC, and A_n , and the core parameters are SLA and N_{area} (Onoda et al., 2017; Yang et al., 2019). PC and A_n are also key to photosynthetic acclimation (Smith and Keenan, 2020; Tcherkez et al., 2020), and A_n is affected by photosynthetic limitations. SLA or leaf mass per unit leaf area ($1/SLA$) is related to leaf thickness and leaf mass density (Maenpuen et al., 2022), reflecting the construction cost of PC (Onoda et al., 2017). N_{area} reflects the nitrogen used for photosynthesis and growth (structure and metabolism) (Krämer et al., 2022). As nitrogen related to photosynthesis usually accounts for about 50% of the total nitrogen content in leaves, there




FIGURE 3 The relative limitations of photosynthesis by biochemistry (l_b), and mesophyll resistance (l_m), and stomatal resistance (l_s) in response to CO_2 and N supply in wintersweet. N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. Abbreviations are provided in Table S1.




FIGURE 4 The A/C_i curves in response to different N levels wintersweet seedlings grown under ambient CO_2 (aCO_2) and elevated CO_2 (eCO_2). N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. Each point denotes the means of six (C_i , A_n) values in A/C_i curve. The circles represent the transition point (C_{i-t} , A_{n-t}) from Rubisco carboxylation to RuBP regeneration of photosynthesis limitation. The triangles indicated the photosynthetic rate (C_{i-g} , A_{n-g}) under growth C_a ($400 \mu\text{mol mol}^{-1}$ versus $800 \mu\text{mol mol}^{-1}$). ACE: apparent carboxylation efficiency estimated from the initial slope of A/C_i curve. Different letters of the same parameter in the same species are significantly different between treatments (Tukey host hoc test, $P < 0.05$, see Table S2).

TABLE 2 Photosynthetic capacity, chlorophyll concentration, conductance to CO_2 diffusion, and intercellular to external CO_2 concentration ratio in wintersweet grown at ambient or elevated CO_2 in response to different N levels.

CO_2	N	V_{cmax} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	J_{max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	Chl (mg m^{-2})	g_m ($\text{mol m}^{-2} \text{s}^{-1}$)	g_s ($\text{mmol m}^{-2} \text{s}^{-1}$)	C_i/C_a
a CO_2	N3	39.5 ± 1 a	63.4 ± 2 ab	350.4 ± 16.5 a	0.21 ± 0.011 ab	42 ± 5.9 a	0.43 ± 0.02 b
	N6	41.9 ± 1.5 a	64.3 ± 2.9 ab	344.4 ± 11.2 a	0.22 ± 0.003 a	37.1 ± 2.3 ab	0.41 ± 0.01 b
	N9	41.2 ± 1 a	69.8 ± 3.3 ab	338.1 ± 10.4 a	0.24 ± 0.009 a	36 ± 3.9 ab	0.42 ± 0.03 b
e CO_2	N3	25.8 ± 2.8 c	54.7 ± 3.4 b	300.2 ± 29.9 a	0.17 ± 0.012 bc	26.2 ± 3.9 b	0.53 ± 0.04 a
	N6	27.3 ± 2.6 bc	60.5 ± 5.2 ab	315.2 ± 27.7 a	0.16 ± 0.011 c	29.2 ± 6 ab	0.6 ± 0.05 a
	N9	32.3 ± 3.1 b	73.7 ± 4.6 a	366.7 ± 31.2 a	0.21 ± 0.017 ab	31.2 ± 3.7 ab	0.56 ± 0.03 a
P-value	CO_2	< 0.001	0.34	0.37	< 0.001	0.01	< 0.001
	N	0.18	< 0.001	0.46	0.01	0.98	0.82
	$\text{CO}_2 \times \text{N}$	0.38	0.25	0.22	0.28	0.46	0.49

Note: each value represents mean \pm SE ($n = 6$). Two-way ANOVA was performed on the effects of CO_2 , nitrogen form (N), and their interactions ($\text{CO}_2 \times \text{N}$). Significant effects ($P \leq 0.05$) are marked in bold and “*” indicates interaction. Different letters within the same column indicated statistically significant differences between treatments (Tukey post hoc test, $P < 0.05$). a CO_2 : ambient CO_2 ($400 \mu\text{mol mol}^{-1}$); e CO_2 : elevated CO_2 ($800 \mu\text{mol mol}^{-1}$); N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. V_{cmax} : maximum rate of ribulose-1,5-bisphosphate carboxylation; J_{max} : maximum photosynthetic electron transport rate; Chl: leaf chlorophyll concentration. g_m : mesophyll conductance; g_s : stomatal conductance; C_i/C_a : intercellular (C_i) to ambient (C_a) CO_2 concentration ratio.

is a trade-off in nitrogen allocation between photosynthesis and growth (Silva-Pérez et al., 2020). Therefore, investigating changes in SLA and N_{area} under different environments is a classic methodology for analyzing plant functions by LES (Onoda et al., 2017; Xiong and Flexas, 2018). At one end of the LES are higher SLA and N_{area} , indicating that a given investment in leaf biomass will lead to a larger light-intercepting area and, higher returns in carbohydrates produced and faster growth. These traits are associated with invasive plant species

(Onoda et al., 2017; Cui et al., 2020). The other end of the LES is associated with higher leaf structure investments, lower returns in carbohydrate production and, slower growth but longer leaf lifespan and robust stress tolerance (Xiong and Flexas, 2018). However, the existence of these relationships depends on sufficient resources (Onoda et al., 2017). Our results showed no synchronous change in SLA and N_{area} , and elevated CO_2 significantly reduced SLA but increased N_{area} ,




FIGURE 5 Principal Component Analysis (PCA) on growth and photosynthetic parameters in wintersweet exposed to CO₂ (a) and N levels (b) treatments. The arrows point to near overlap, vertical, and reverse, which represent positive, no, and negative correlations between these parameters respectively. N3, N6 and N9 mean that fertilized 3 mM, 6 mM and 9 mM nitrogen respectively. A_{n-t} : net photosynthesis rate at transition point (C_{i-t} , A_{n-t}) between Rubisco limitation and RuBP regeneration limitation based on A/C_i curve; N_{area} : leaf N per unit area; C_i/C_a : the ratio of C_i and C_a ; A_{n-g} : net photosynthesis rate at a growth [CO₂] which eCO₂ at 800 $\mu\text{mol mol}^{-1}$ and aCO₂ at 400 $\mu\text{mol mol}^{-1}$; C/N: leaf carbon and nitrogen ratio; A_{max} : photosynthetic rate of saturation light at 400 $\mu\text{mol mol}^{-1}$; L_s : stomatal limitation; $iWUE$: intrinsic water use efficiency; SLA : specific leaf area; $PNUE$: photosynthesis nitrogen use efficiency; ACE : apparent carboxylation efficiency estimated from the initial slope of A/C_i curve; V_{cmax} : maximum rate of ribulose-1,5-bisphosphate carboxylation; g_m : mesophyll conductance; g_s : stomatal conductance; Chl : leaf chlorophyll concentration; J_{max} : maximum of photosynthetic electron transport rate. See Table S1 for other explanations.

suggesting that elevated CO₂ might promote wintersweet's growth, lifespan, and stress resistance.

Elevated CO₂ generally improves carbon fixation and biomass production, diluting leaf nitrogen concentration and reducing photosynthetic capacity (Ainsworth and Long, 2021). However, our results show that eCO₂ reduced PC and chlorophyll concentration but increased N_{area} , indicating a reduction of N allocation to photosynthetic apparatus, particularly light-harvesting pigments (N_v). It is unclear where the nitrogen saved from the reduction of the allocation to photosynthetic apparatus was allocated. However, this change in N allocation did reduce the photosynthetic nitrogen use efficiency in seedlings grown under eCO₂, although eCO₂ significantly increased A_{n-g} . $PNUE$ is closely related to the proportion of leaf N allocated to photosynthesis (mainly to Rubisco) (Onoda et al., 2017). Our principal component analysis and deconstructed equation model results suggest that increasing nitrogen supply appears to have compensated for the negative effects of elevated CO₂ on photosynthetic capacity. Therefore, the elevated CO₂ and higher nitrogen supply synergistically affected the growth of Wintersweet.

4.2 | Acclimation to elevated CO₂ and biochemical limitations to photosynthesis

Elevated CO₂ usually decreases V_{cmax} and J_{max} , a phenomenon known as photosynthetic acclimation or downregulation (Smith and Keenan, 2020; Tcherkez et al., 2020). CO₂ is a substrate for photosynthesis, and a higher concentration will reduce the substrate limitation for photosynthesis, leading to a decrease in N allocation to photosynthetic machinery (Silva-Pérez et al., 2020). We observed that

Wintersweet exhibited a typical photosynthetic acclimation reaction. The effect of photosynthetic downregulation, however, is generally smaller than the stimulation of photosynthesis by the elevation of CO₂, resulting in a higher actual rate of A_{n-g} under eCO₂ than under aCO₂ (Wang et al., 2022), as we have observed in this current study.

The eCO₂ did not change the primary limitation of photosynthesis in wintersweet. Some previous studies have reported that the elevation of CO₂ can trigger the shift of the primary limitation of photosynthesis from biochemical limitation (i.e., V_{cmax} limitation, $A_{n-g} < A_{n-t}$) to photochemical limitation (i.e., limitation by J_{max} , $A_{n-g} > A_{n-t}$) (Wang et al., 2023). However, the photosynthesis of wintersweet was still primarily limited by V_{cmax} under eCO₂ in this study (Figure 4). Furthermore, the eCO₂ significantly reduced leaf chlorophyll concentration in wintersweet, which is generally positively correlated to light harvesting and photosynthetic electron transport. These results suggest that the resistance to CO₂ diffusion from the atmosphere to the carboxylation site in the chloroplast will likely continue to exert a major limitation to the photosynthesis of wintersweet in the future with elevated CO₂.

4.3 | CO₂ diffusion limitations to photosynthesis

An interesting finding of this study is that wintersweet had very low stomatal conductance, less than 50 $\text{mmol m}^{-2} \text{s}^{-1}$ in all treatments, which is much lower than typical values for deciduous broad-leaved tree species under non-stress conditions (Wang et al., 2022). In contrast, the mesophyll conductance values, g_m , were comparable to those reported for other deciduous tree species (Elferjani et al., 2021). For most tree species, the values of g_s and g_m are about equal (Wang and Dang, 2022). The g_s of wintersweet, however, was only about 1/6




FIGURE 6 The *priori* (showing all tested paths) piecewise structural equation model (pSEM) relating to direct and indirect effects of CO₂ and N levels on biomass and A_{n-g} . Two CO₂ treatments (ambient CO₂: 400 $\mu\text{mol mol}^{-1}$ and elevated CO₂: 800 $\mu\text{mol mol}^{-1}$) and N levels (fertilized 3 mM (N3), 6 mM (N6) and 9 mM (N9) nitrogen) are treated as categorical variables. A_{n-g} : net photosynthesis rate at a growth [CO₂] which eCO₂ at 800 $\mu\text{mol mol}^{-1}$ and aCO₂ at 400 $\mu\text{mol mol}^{-1}$; N_{area} : leaf N per unit area; SLA: specific leaf area; *Chl*: leaf chlorophyll concentration; g_t : total conductance to CO₂ between the leaf surface and carboxylation sites ($1/g_t = 1/g_s + 1/g_m$). Multiple regression from V_{cmax} and J_{max} to A_{n-g} was used to construct photosynthetic capacity (PC) as a component variable (hexagonal frame). The SEM (using psem function in piecewiseSEM package R) of biomass and A_{n-g} in wintersweet by biomass (a) and by A_{n-g} (b). Arrows mean the directional influence between the variables and the solid lines represent significant relationships, and the dashed lines refer to nonsignificant relationships ($P > 0.05$). The numbers near the arrows represent the standardized path coefficients (for continuous variables), and the numbers under the box with R^2 refer to the degree of variation of the variable interpreted by all paths. Curved double arrows represent a significant correlation between variables (with correlation coefficient). The values on the line from the categorical variable indicated the continuous variable estimated marginal means by the treatment levels.

of g_m and more than 50% of the limitation to photosynthesis was attributable to stomatal conductance, which is very high compared to other tree species (Sakoda et al., 2021). The results of the relative

limitation analysis of photosynthesis further corroborated this conclusion. Furthermore, the results of the structural equation model indicate that CO₂ conductivity had a direct effect on A_{n-g} , and the influence coefficient was 0.61 high, further confirming that the photosynthesis of Wintersweet was mainly limited by the resistance to CO₂ diffusion (Figure 6). Elevated CO₂ further increased the proportion of stomatal limitation to photosynthesis in wintersweet. The results suggest that wintersweet is a very conservative species for water use and its water use efficiency will likely be further enhanced by elevated CO₂ in the future. The high-water use efficiency is a highly desirable trait for landscaping trees.

5 | CONCLUSIONS

As a popular and special landscape tree species, wintersweet plays an important role in urban landscapes, particularly for the winter season. The unique fresh floral scent of its yellow flowers makes people feel magical and quiet in the cold winter. Its high drought resistance may be related to its low g_s . A lower g_s has a greater limitation to transpiration than to photosynthesis, enhancing water use efficiency. However, the increase in leaf area (canopy) under elevated CO₂ indicates that the species may have increased growth in the future. The LES with elevated CO₂ showed a decrease in SLA and an increase in N_{area} , indicating that future climate conditions may be favourable for the growth and stress resistance of wintersweet. Wintersweet exhibited typical symptoms of photosynthetic acclimation, such as decreased PC in response to elevated CO₂. However, leaf nitrogen concentration increased, indicating that Wintersweet may be more sensitive to future climate change and soil nitrogen levels.

We proposed and tested the framework to use a combination of LES and photosynthetic acclimation to evaluate the potential performance of Wintersweet under future CO₂. The results suggest that wintersweet should be cultivated more in future urban landscape planning, and nitrogen fertilization should further enhance growth and physiological performance under the predicted future CO₂ levels. Our results suggest that elevated CO₂ can further enhance the water conservation and water use efficiency of wintersweet in the future, making it a highly desirable species for landscaping in the future when more frequent and more severe drought spells are predicted under the scenario of climate change.

AUTHOR CONTRIBUTIONS

Lei Wang: Conceptualization, Methodology, Data curation, Writing – original draft. Qing-lai DANG: Conceptualization, Writing – review & editing.

ACKNOWLEDGEMENTS

We want to thank Ms. Keri Pidgen, the Greenhouse Manager, for her logistic support and other operational assistance during the experiments. The study was supported by NSERC Discovery Development Grant to QL Dang (Project No. DDG-2020-00008) and by Zhejiang

A&U University Jiyang College Talent Initiation Project to Lei Wang (RC2023B06).

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no known competing financial interests of personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article.

ORCID

Qing-Lai Dang  <https://orcid.org/0000-0002-5930-248X>

REFERENCES

- Ainsworth, E.A., Long, S.P., 2021. 30 years of free-air carbon dioxide enrichment (FACE): What have we learned about future crop productivity and its potential for adaptation? *Global change biology* 27, 27–49.
- Alizadeh, B., Hitchmough, J.D., 2020. How will climate change affect future urban naturalistic herbaceous planting? The role of plant origin and fitness. *Urban Forestry & Urban Greening* 54, 126786.
- Anirban, B., 2019. Finding the fragrance genes of wintersweet. *Physiologia plantarum* 166, 475–477.
- Bae, J., Ryu, Y., 2015. Land use and land cover changes explain spatial and temporal variations of the soil organic carbon stocks in a constructed urban park. *Landscape and Urban Planning* 136, 57–67.
- Bernacchi, C.J., Singsaas, E.L., Pimentel, C., Portis Jr, A.R., Long, S.P., 2001. Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, cell & environment* 24, 253–259.
- Bowen, J.L., Kearns, P.J., Byrnes, J.E.K., Wigginton, S., Allen, W.J., Greenwood, M., Tran, K., Yu, J., Cronin, J.T., Meyerson, L.A., 2017. Lineage overwhelms environmental conditions in determining rhizosphere bacterial community structure in a cosmopolitan invasive plant. *Nature communications* 8, 433.
- Cui, E., Weng, E., Yan, E., Xia, J., 2020. Robust leaf trait relationships across species under global environmental changes. *Nature communications* 11, 2999.
- Dervishi, V., Fleckenstein, C., Rahman, M.A., Pauleit, S., Ludwig, F., Pretzsch, H., Rötzer, T., 2023. Trees in planters – Growth, structure and ecosystem services of *Platanus x hispanica* and *Tilia cordata* and their reaction to soil drought. *Urban Forestry & Urban Greening* 86, 128024.
- Dowtin, A.L., Cregg, B.C., Nowak, D.J., Levia, D.F., 2023. Towards optimized runoff reduction by urban tree cover: A review of key physical tree traits, site conditions, and management strategies. *Landscape and Urban Planning* 239, 104849.
- Dujardin, S., Stas, M., van Eupen, C., Aerts, R., Hendrickx, M., Delclocq, A.W., Duchêne, F., Hamdi, R., Nawrot, T.S., van Nieuwenhuyse, Aerts, J.-M., van Orshoven, J., Somers, B., Linard, C., Dendoncker, N., 2022. Mapping abundance distributions of allergenic tree species in urbanized landscapes: A nation-wide study for Belgium using forest inventory and citizen science data. *Landscape and Urban Planning* 218, 104286.
- Duursma, R.A., 2015. Plantcophys--An R Package for Analysing and Modelling Leaf Gas Exchange Data. *PLoS one* 10, e0143346.
- Elferjani, R., Benomar, L., Momayyezi, M., Tognetti, R., Niinemets, Ü., Soolanayakanahally, R.Y., Thérroux-Rancourt, G., Tosens, T., Ripullone, F., Bilodeau-Gauthier, S., Lamhamedi, M.S., Calfapietra, C., Lamara, M., 2021. A meta-analysis of mesophyll conductance to CO₂ in relation to major abiotic stresses in poplar species. *Journal of experimental botany* 72, 4384–4400.
- Esperon-Rodriguez, M., Quintans, D., Rymer, P.D., 2023. Urban tree inventories as a tool to assess tree growth and failure: The case for Australian cities. *Landscape and Urban Planning* 233, 104705.
- Falster, D.S., Duursma, R.A., FitzJohn, R.G., 2018. How functional traits influence plant growth and shade tolerance across the life cycle. *Proceedings of the National Academy of Sciences of the United States of America* 115, E6789–E6798.
- Farquhar, J.A., Caemmerer, S., Berry, J.A., 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149, 78–90.
- Firn, J., McGree, J.M., Harvey, E., Flores-Moreno, H., Schütz, M., Buckley, Y.M., Borer, E.T., Seabloom, E.W., La Pierre, K.J., MacDougall, A.M., Prober, S.M., Stevens, C.J., Sullivan, L.L., Porter, E., Ladouceur, E., Allen, C., Moromizato, K.H., Morgan, J.W., Harpole, W.S., Hautier, Y., Eisenhauer, N., Wright, J.P., Adler, P.B., Arnillas, C.A., Bakker, J.D., Biederman, L., Broadbent, A.A.D., Brown, C.S., Bugalho, M.N., Caldeira, M.C., Cleland, E.E., Ebeling, A., Fay, P.A., Hagenah, N., Kleinhesselink, A.R., Mitchell, R., Moore, J.L., Nogueira, C., Peri, P.L., Roscher, C., Smith, M.D., Wragg, P.D., Risch, A.C., 2019. Leaf nutrients, not specific leaf area, are consistent indicators of elevated nutrient inputs. *Nature ecology & evolution* 3, 400–406.
- Gratani, L., Varone, L., 2007a. Plant crown traits and carbon sequestration capability by *Platanus hybrida* Brot. in Rome. *Landscape and Urban Planning* 81, 282–286.
- Gratani, L., Varone, L., 2007b. Plant crown traits and carbon sequestration capability by *Platanus hybrida* Brot. in Rome. *Landscape and Urban Planning* 81, 282–286.
- IPCC, 2022. *Climate Change 2022 - Mitigation of Climate Change - Full Report*. Cambridge University Press.
- Killi, D., Bussotti, F., Gottardini, E., Pollastrini, M., Mori, J., Tani, C., Papini, A., Ferrini, F., Fini, A., 2018. Photosynthetic and morphological responses of oak species to temperature and [CO₂] increased to levels predicted for 2050. *Urban Forestry & Urban Greening* 31, 26–37.
- Kim, J., Lee, D.K., Kim, H.G., 2020. Suitable trees for urban landscapes in the Republic of Korea under climate change. *Landscape and Urban Planning* 204, 103937.
- Kimm, H., Ryu, Y., 2015. Seasonal variations in photosynthetic parameters and leaf area index in an urban park. *Urban Forestry & Urban Greening* 14, 1059–1067.
- Knauer, J., Zaehle, S., Kauwe, M.G. de, Bahar, N.H.A., Evans, J.R., Medlyn, B.E., Reichstein, M., Werner, C., 2019. Effects of mesophyll conductance on vegetation responses to elevated CO₂ concentrations in a land surface model. *Global change biology* 25, 1820–1838.
- Krämer, K., Kepp, G., Brock, J., Stutz, S., Heyer, A.G., 2022. Acclimation to elevated CO₂ affects the C/N balance by reducing de novo N-assimilation. *Physiologia plantarum* 174, e13615.
- Lefcheck, J.S., 2016. piecewiseSEM: Piecewise structural equation modeling in R for ecology, evolution, and systematics. *Methods Ecol Evol* 7, 573–579.
- Liu, H., Ye, Q., Simpson, K.J., Cui, E., Xia, J., 2022. Can evolutionary history predict plant plastic responses to climate change? *The New phytologist* 235, 1260–1271.
- Liu, J., Slik, F., 2022. Are street trees friendly to biodiversity? *Landscape and Urban Planning* 218, 104304.
- Lo Piccolo, E., Lauria, G., Pellegrini, E., Cotrozzi, L., Guidi, L., Skoet, M., Vernieri, P., Remorini, D., Massai, R., Landi, M., 2023. Testing the suitability for coastal green areas of three ornamental shrub species through physiological responses to the saline nebulization. *Urban Forestry & Urban Greening* 84, 127920.
- Lu, Z., Ren, T., Li, J., Hu, W., Zhang, J., Yan, J., Li, X., Cong, R., Guo, S., Lu, J., 2020. Nutrition-mediated cell and tissue-level anatomy triggers the covariation of leaf photosynthesis and leaf mass per area. *Journal of experimental botany* 71, 6524–6537.
- Maenpuen, P., Katabuchi, M., Onoda, Y., Zhou, C., Zhang, J.-L., Chen, Y.-J., 2022. Sources and consequences of mismatch between leaf disc and

- whole-leaf leaf mass per area (LMA). *American journal of botany* 109, 1242–1250.
- Midolo, G., Frenne, P. de, Hölzel, N., Wellstein, C., 2019. Global patterns of intraspecific leaf trait responses to elevation. *Global change biology* 25, 2485–2498.
- Nitschke, C.R., Nichols, S., Allen, K., Dobbs, C., Livesley, S.J., Baker, P.J., Lynch, Y., 2017. The influence of climate and drought on urban tree growth in southeast Australia and the implications for future growth under climate change. *Landscape and Urban Planning* 167, 275–287.
- Nyelele, C., Kroll, C.N., 2021. A multi-objective decision support framework to prioritize tree planting locations in urban areas. *Landscape and Urban Planning* 214, 104172.
- Onoda, Y., Wright, I.J., Evans, J.R., Hikosaka, K., Kitajima, K., Niinemets, Ü., Poorter, H., Tosens, T., Westoby, M., 2017. Physiological and structural tradeoffs underlying the leaf economics spectrum. *The New phytologist* 214, 1447–1463.
- Ordóñez Barona, C., Wolf, K., Kowalski, J.M., Kendal, D., Byrne, J.A., Conway, T.M., 2022. Diversity in public perceptions of urban forests and urban trees: A critical review. *Landscape and Urban Planning* 226, 104466.
- Pao, Y.-C., Chen, T.-W., Moualeu-Ngangue, D.P., Stützel, H., 2019. Environmental triggers for photosynthetic protein turnover determine the optimal nitrogen distribution and partitioning in the canopy. *Journal of experimental botany* 70, 2419–2434.
- Sakoda, K., Yamori, W., Groszmann, M., Evans, J.R., 2021. Stomatal, mesophyll conductance, and biochemical limitations to photosynthesis during induction. *Plant physiology* 185, 146–160.
- Shang, J., Tian, J., Cheng, H., Yan, Q., Li, L., Jamal, A., Xu, Z., Xiang, L., Sasaki, C.A., Jin, S., Zhao, K., Liu, X., Chen, L., 2020. The chromosome-level wintersweet (*Chimonanthus praecox*) genome provides insights into floral scent biosynthesis and flowering in winter. *Genome biology* 21, 200.
- Shen, Z., Li, W., Li, Y., Liu, M., Cao, H., Provar, N., Ding, X., Sun, M., Tang, Z., Yue, C., Cao, Y., Yuan, D., Zhang, L., 2021. The red flower wintersweet genome provides insights into the evolution of magnoliids and the molecular mechanism for tepal color development. *The Plant journal: for cell and molecular biology* 108, 1662–1678.
- Silva-Pérez, V., Faveri, J. de, Molero, G., Deery, D.M., Condon, A.G., Reynolds, M.P., Evans, J.R., Furbank, R.T., 2020. Genetic variation for photosynthetic capacity and efficiency in spring wheat. *Journal of experimental botany* 71, 2299–2311.
- Simon, H., Lindén, J., Hoffmann, D., Braun, P., Bruse, M., Esper, J., 2018. Modeling transpiration and leaf temperature of urban trees – A case study evaluating the microclimate model ENVI-met against measurement data. *Landscape and Urban Planning* 174, 33–40.
- Smith, N.G., Keenan, T.F., 2020. Mechanisms underlying leaf photosynthetic acclimation to warming and elevated CO₂ as inferred from least-cost optimality theory. *Global change biology* 26, 5202–5216.
- Song, X., Wu, Q., 2022. Study on smellscape perception and landscape application of fragrant plants. *Urban Forestry & Urban Greening* 67, 127429.
- Sousa-Silva, R., Duflos, M., Ordóñez Barona, C., Paquette, A., 2023. Keys to better planning and integrating urban tree planting initiatives. *Landscape and Urban Planning* 231, 104649.
- Stas, M., Aerts, R., Hendrickx, M., Dendoncker, N., Dujardin, S., Linard, C., Nawrot, T., van Nieuwenhuysse, Aerts, J.-M., van Orshoven, J., Somers, B., 2020. An evaluation of species distribution models to estimate tree diversity at genus level in a heterogeneous urban-rural landscape. *Landscape and Urban Planning* 198, 103770.
- Tang, L., Li, D., Ma, Y., Wang, J., Tian, Y., 2023. The association of residential greenness, genetic susceptibility and telomere length: A cross-sectional study. *Landscape and Urban Planning* 239, 104864.
- Tcherkez, G., Ben Mariem, S., Larraya, L., García-Mina, J.M., Zamarreño, A.M., Paradela, A., Cui, J., Badeck, F.-W., Meza, D., Rizza, F., Bunce, J., Han, X., Tausz-Posch, S., Cattivelli, L., Fangmeier, A., Aranjuelo, I., 2020. Elevated CO₂ has concurrent effects on leaf and grain metabolism but minimal effects on yield in wheat. *Journal of experimental botany* 71, 5990–6003.
- Tian, J.-P., Ma, Z.-Y., Zhao, K.-G., Zhang, J., Xiang, L., Chen, L.-Q., 2019a. Transcriptomic and proteomic approaches to explore the differences in monoterpene and benzenoid biosynthesis between scented and unscented genotypes of wintersweet. *Physiologia plantarum* 166, 478–493.
- Tian, J.-P., Ma, Z.-Y., Zhao, K.-G., Zhang, J., Xiang, L., Chen, L.-Q., 2019b. Transcriptomic and proteomic approaches to explore the differences in monoterpene and benzenoid biosynthesis between scented and unscented genotypes of wintersweet. *Physiologia plantarum* 166, 478–493.
- Trouwborst, G., Hogewoning, S.W., Harbinson, J., van Ieperen, W., 2011. Photosynthetic acclimation in relation to nitrogen allocation in cucumber leaves in response to changes in irradiance. *Physiologia plantarum* 142, 157–169.
- Velasco, E., Roth, M., Norford, L., Molina, L.T., 2016. Does urban vegetation enhance carbon sequestration? *Landscape and Urban Planning* 148, 99–107.
- Walker, B.J., Skabelund, D.C., Busch, F.A., Ort, D.R., 2016. An improved approach for measuring the impact of multiple CO₂ conductances on the apparent photorespiratory CO₂ compensation point through slope-intercept regression. *Plant, cell & environment* 39, 1198–1203.
- Wang, L., Dang, Q.-L., 2022. Growth and photosynthetic traits differ between shoots originated from axillary buds or from adventitious buds in *Populus balsamifera* L. cuttings. *Physiologia plantarum* 174, e13599.
- Wang, L., Zheng, J., Wang, G., Dang, Q.-L., 2022. Increased leaf area compensated photosynthetic downregulation in response to elevated CO₂ and warming in white birch. *Can. J. For. Res.* 52, 1176–1185.
- Wang, L., Zheng, J., Wang, G., Dang, Q.-L., 2023. Combined effects of elevated CO₂ and warmer temperature on limitations to photosynthesis and carbon sequestration in yellow birch. *Tree physiology* 43, 379–389.
- Wang, Y., Chang, Q., Li, X., 2021. Promoting sustainable carbon sequestration of plants in urban greenspace by planting design: A case study in parks of Beijing. *Urban Forestry & Urban Greening* 64, 127291.
- Wellburn, A., 1994. The Spectral Determination of Chlorophylls a and b, as well as Total Carotenoids, Using Various Solvents with Spectrophotometers of Different Resolution. *Journal of plant physiology* 144, 307–313.
- Xiong, D., Flexas, J., 2018. Leaf economics spectrum in rice: leaf anatomical, biochemical, and physiological trait trade-offs. *Journal of experimental botany* 69, 5599–5609.
- Yang, Y., Wang, H., Harrison, S.P., Prentice, I.C., Wright, I.J., Peng, C., Lin, G., 2019. Quantifying leaf-trait covariation and its controls across climates and biomes. *The New phytologist* 221, 155–168.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Wang, L. & Dang, Q.-L. (2024) Using leaf economic spectrum and photosynthetic acclimation to evaluate the potential performance of wintersweet under future climate conditions. *Physiologia Plantarum*, 176(3), e14318. Available from: <https://doi.org/10.1111/ppl.14318>